轻中度镇静-口服或综合笑气镇静剂的术前和术后指南

为了病人的安全，请详细阅读并遵从下列重要指南

饮食 为了避免在治疗期间出现呕吐及其它并发症，请在治疗前六小时禁食，**不要**给孩子吃任何食物或喝饮料。

健康状况变化 若孩子在治疗前三天内有感冒，剧烈咳嗽或发烧，请尽快与我们联系重新预约。 在治疗当日也请在治疗前告知诊所工作人员有关孩子健康状况的任何变化，。

赴约 您的孩子去诊所就诊时需要监护人陪同前往。该监护人需要留在诊所等候区直到治疗结束。

药物 在治疗前后，请只给孩子服用每天必吃的药物，比如癫痫药物或预防性抗生素以及孩子的家庭医生开的处方药。请告知诊所人员需服用的药物。除此以外的其它任何药物，请务必与诊所确认后才能给孩子服用。

活动 在治疗后，孩子需有人密切观察五个小时以防有超过30分钟的恶心不适感。 **孩子当天不能去上学，也要避免任何活动，需要在家休息。**

回家 您的孩子必须由监护人陪伴并由专人驾车护送回家。 在乘车时请按规定使用孩童座椅和安全带。途中监护人需要密切观察孩子有无呼吸困难等情况

治疗后的饮食 治疗结束后为孩子补充水分非常重要。第一次补充只需要水，之后可以给一些无渣的甜饮料。在孩子想进食时，请以多次少量的方式给孩子凉的软性食物。

面颊 / 嘴唇和舌头的麻木感

治疗后孩子的面颊，嘴唇和舌头可能出现麻木感。请密切观察孩子，确保他们不要咬到自己的面颊，嘴唇或舌头。

疼痛缓解与护理 每 6 - 8 小时给孩子吃一次儿童型对乙酰氨基酚 (比如泰诺Tylenol)或布洛芬 (比如 Advil或者 Motrin) 帮助缓解疼痛和不适。**如有以下情况，请马上与我们电话联系**：

1. 呕吐持续四个小时以上
2. 体温持续升高长达二十四小时以上，或者体温高于 101F (38度 C)
3. 呼吸困难
4. 其他任何您担心的情况。

如果您有任何与治疗有关的疑虑，请与我们联系。诊所下班后的紧急呼机号码为 604-734-7243

Certified Specialists in Pediatric Dentistry

Anabel R. Chan, DDS, Dip Peds, MSc, FRCDC
M-Reza Nouri DMD, Dip Peds, MSc, FRCDC
Louisa Y. Leung, DMD, Cert. Peds, MS, FRCDC
Carter K. Ng, DDS, Dip Peds, MSc, FRCDC
Bradford W. Scheideman, DMD, Cert. Peds, FRCDC
& Associates

Certified Specialists in Orthodontics

Donal C. Flanagan, DDS, MS, FRCDC
Christian A. Wong, DDS, MS, FRCDC
Todd R. Moore, DDS, MS, FRCDC
& Associates
PDG Pediatric Dentistry & Orthodontics
Vancouver - Richmond - Delta/Surrey – Coquitlam
www.PDGdental.com 604-734-1000